Mon. Not. R. Astron. Soc. **000**, 000–000 (0000) Printed 5 February 2008 (MN LaTeX style file v1.4)

arXiv:astro-ph/9911513v3 15 Aug 2000

Influence of the r-mode instability on hypercritically accreting neutron stars

Shin'ichirou Yoshida ⋆ † and Yoshiharu Eriguchi‡

Department of Earth Science and Astronomy, Graduate School of Arts and Sciences, University of Tokyo,
Komaba, Meguro-ku, Tokyo 153-8902, Japan

Accepted, Received

ABSTRACT

We have investigated an influence of the r-mode instability on hypercritically accreting ($\dot{M} \sim 1 M_\odot \mathrm{y}^{-1}$) neutron stars in close binary systems during their common envelope phases based on the scenario proposed by Bethe et al. (1999). On the one hand neutron stars are heated by the accreted matter at the stellar surface, but on the other hand they are also cooled down by the neutrino radiation. At the same time, the accreted matter transports its angular momentum and mass to the star. We have studied the evolution of the stellar mass, temperature and rotational frequency.

The gravitational-wave-driven instability of the r-mode oscillation strongly suppresses spinning-up of the star, whose final rotational frequency is well below the mass-shedding limit, typically as small as 10% of that of the mass-shedding state. On a very short time scale the rotational frequency tends to approach a certain constant value and saturates there as far as the amount of the accreted mass does not exceed a certain limit to collapse to a black hole. This implies that the similar mechanism of gravitational radiation as the so-called Wagoner star may work in this process. The star is spun up by accretion until the angular momentum loss by gravitational radiation balances the accretion torque. The time-integrated dimensionless strain of the radiated gravitational wave may be large enough to be detectable by the gravitational wave detectors such as LIGO II.

Key words: accretion – binaries: close – stars: neutron – stars: rotation – stars: oscillations

1 INTRODUCTION

In relation to the formation mechanism of compact binary systems such as double neutron star systems, double black hole systems or black hole–neutron star systems, hypercritical accretion flows onto neutron stars in envelopes of massive stars of close binary systems have been investigated by several authors (Chevalier 1993; Chevalier 1996; Brown 1995; Bethe & Brown 1998; Bethe et al. 1999). The mass accretion rate ($\dot{M}$) of the hypercritical accretion flow amounts to 10^8 times as high as the Eddington's mass accretion limit. Chevalier (1996) found that for rather small values of the viscosity parameter, $\alpha \sim 10^{-6}$, the accretion flow which cools via neutrino radiation could be realized.

Recently Bethe et al. (1999) pointed out that the flow with much larger and conservative values of the viscosity parameter, $\alpha \gtrsim 0.1$, does not become so hot as to emit neutrinos during the accretion process but that the accretion will proceed to the advection dominated accretion flow (ADAF). In their model, the accreted matter onto the neutron star surface is processed there by nuclear reactions and cools via neutrino emission. The hypercritical accretion increases the mass of the star rapidly and induces the collapse of the neutron star to a black hole. Thus the supernova explosion of the massive companion will lead to the formation of a black hole–neutron star binary system or a double black hole binary system. The merging event of such a system might be one of the important targets for the gravitational wave astronomy.

In the accretion process, a non-negligible amount of angular momentum is carried into the neutron star. This potentially could spin up the star to the Kepler limit, i.e. to the state where the stellar rotational frequency equals to the orbital frequency at the equatorial surface. The star rotating at the Kepler limit deforms significantly from the spherical configuration and the subsequent collapse of the rapidly

---

⋆ yoshida@sissa.it

† present address: Scuola Internazionale Superiore di Studi Avanzati, Via Beirut 2-4, 34014 Trieste, Italy

‡ eriguchi@valis.c.u-tokyo.ac.jp

© 0000 RAS

rotating star to a black hole might radiate a substantial fraction of the binding energy of the system in the form of gravitational waves.

This seemingly straightforward conclusion, however, has to be reexamined from the standpoint of stability of rotating neutron stars. The gravitational radiation driven instability of rotating stars might prevent the stellar rotational frequency to increase above a certain limit. In particular, the recently discovered r-mode instability (Andersson 1998; Friedman & Morsink 1998) might be sufficiently strong to suppress the spin frequency in some range of the stellar temperature.

In this paper, in order to see how this instability influences the accretion induced spin-up of the neutron star in the hypercritical accretion, we will employ the hypercritical ADAF model based on the analysis of Bethe et al. (1999) and investigate the evolution of the temperature and the rotational frequency of the neutron star. §

2 SIMPLIFIED MODEL OF THE ACCRETION INDUCED EVOLUTION OF NEUTRON STARS

It should be noted that the study in this paper is done in the framework of Newtonian dynamics and Newtonian gravity. The gravitational radiation reaction on stellar oscillations is evaluated by the lowest order post-Newtonian approximation (Thorne 1980). Furthermore to simplify our analysis, stellar configurations are assumed to be spherical polytropes, $p = K\rho^2$, with the polytropic index $N = 1$. Here p and ρ are the pressure and the density, respectively. By choosing the polytropic coefficient K appropriately, the stellar mass is $M = 1.4M_\odot$ and its radius is $R = 12.5$km. Note that the radius of the polytropes with index $N = 1$ is independent of the central density and thus of the mass if the constant K is fixed.

2.1 Characteristics of the hypercritical ADAF onto neutron stars

The hypercritical accretion flow which we employ in this paper is proposed by Bethe et al. (1999). The characteristics of that flow are summarized as follows. (1) The flow is hypercritical because the accretion rate is typically $\sim 1M_\odot\mathrm{y}^{-1}$. (2) Since the flow is advection dominated (Narayan & Yi 1994) and its temperature is relatively low (~ 0.5MeV), the adiabatic exponent of the gas equals to 4/3 (radiation dominated mixture of photon, electron, positron and nuclei). (3) The accreted matter touches down on the surface of the star 'softly' and burns to produce as heavy nuclei as ^{56}Ni. On the other hand, the matter is cooled by neutrino emission. Consequently the whole process acts as a 'thermostat' which keeps the neutrino temperature $\sim$ 1MeV.

2.2 Mass accretion rate

The mass accretion rate $\dot{M}_{\mathrm{acc}}$, which is the rate of mass attachment to the star, is assumed to be constant during the evolution. In general the mass accretion rate should be distinguished from the mass inflow rate $\dot{M}_{\mathrm{in}}$, which is the total amount of the incoming mass per unit time to the neutron star. This is because some kinds of outflow may coexist with the accreting flows. In this paper we simplify the situation by assuming that both rates are the same, i.e. $\dot{M}_{\mathrm{in}} = \dot{M}_{\mathrm{acc}} \equiv \dot{M}$. We will discuss this assumption further in *Discussion.*

The evolution of the neutron star mass M is written as:

$$\frac{M}{M_\odot} = \mu_0 + \beta t \ , \tag{1}$$

where μ_0 is the initial mass of the star (in units of the solar mass), time t is measured in units of second, and β is defined as:

$$\beta = 3.2 \times 10^{-8} \left[\frac{\dot{M}}{1M_\odot \mathrm{y}^{-1}} \right] \ . \tag{2}$$

2.3 Thermal response of the neutron star heated by neutrino from the surface

The accreted matter landing on the surface layer is processed by nuclear burning and emits neutrinos. We will assume that roughly a half of them escapes from the star freely and that some fraction of the rest of them interacts with stellar matter and heats it up. The mean free path λ of inelastic scattering of neutrinos with electrons in the neutron star matter is written as (Shapiro & Teukolsky 1983),

$$\lambda \sim 2 \times 10^{10} \left[\frac{\rho_{\mathrm{nuc}}}{\bar{\rho}} \right]^{7/6} \left[\frac{0.1\mathrm{MeV}}{E_\nu} \right]^{5/2} \mathrm{cm} \ , \tag{3}$$

where ρ_{nuc} is the 'nuclear density' ($\sim 2.8 \times 10^{14}$g cm^{-3}) and E_ν is the neutrino energy. The density $\bar{\rho}$ means the averaged density of the star, which is related in our polytropic stellar models to the mass M as follows:

$$\frac{\bar{\rho}}{\rho_{\mathrm{nuc}}} = 0.7 \frac{M}{M_\odot} \ . \tag{4}$$

The neutrino energy E_ν is fixed to 1MeV. We simply assume the scattering (thus heating) rate as $\eta R/\lambda$ where R is the stellar radius and η is an efficiency factor of ν-heating through the inelastic scattering. As the flow is adiabatic and radiation dominated, the specific internal energy of the accreted matter ε is written as,

$$\varepsilon = \frac{3}{4} \times \frac{6}{7} \frac{GM}{R} \ . \tag{5}$$

With these expressions, the heating rate of the whole star $\dot{\varepsilon_{\mathrm{h}}}$ is,

$$\begin{aligned} \dot{\varepsilon_{\mathrm{h}}} &\sim \dot{M} \varepsilon \frac{\eta R}{\lambda} \\ &\sim 4.7 \times 10^{43} \left[\frac{\dot{M}}{1M_\odot \mathrm{y}^{-1}} \right] \left[\frac{M}{M_\odot} \right]^{13/6} \mathrm{erg\ s}^{-1}. \end{aligned} \tag{6}$$

Here we choose $\eta = 1$. Heating rate above is proportional to the multiplication of $\dot{M}$ and η. Thus the reduction of η is equivalent to that of $\dot{M}$ with η being fixed. As shown later in Discussion, the dependence of the results on η is weak.[¶]

---

§ Similar investigations of accreting neutron stars with much lower accretion rate, which are the models of low mass X-ray binaries, can be seen for instance in Andersson et al. (1999) and in Levin (1999).

As the cooling mechanism of the star, we assume the modified URCA process to be dominant (Shapiro & Teukolsky 1983). Its cooling rate $L_\nu^{\rm URCA}$ is,

$$L_\nu^{\rm URCA} = 6\times 10^{39}\left[\frac{M}{M_\odot}\right]^{2/3} T_9^8\ , \tag{7}$$

where T_9 is the stellar temperature, which is assumed to be uniform, in units of 10^9K. The heat capacity $C_{\rm v}$ of the neutron star which is approximated by the following formula if the neutron star is assumed to consist of degenerate free nucleons,

$$C_{\rm v} = 8\times 10^{54} k_B \times \left[\frac{M}{M_\odot}\right]\left[\frac{\bar{\rho}}{\rho_{\rm nuc}}\right]^{-2/3} T_9\ , \tag{8}$$

where k_B is the Boltzmann constant. Then the equation of the thermal balance of the star can be written as,

$$\frac{d}{dt}\left(\frac{1}{2}C_{\rm v}T\right) = \dot{\varepsilon}_{\rm h} - L_\nu^{\rm URCA}\ . \tag{9}$$

This is transformed to the equation of the evolution of the temperature as follows:

$$\begin{aligned}\frac{d}{dt}\left(\left[\frac{M}{M_\odot}\right]^{1/3} T_9^2\right) &= 6.5\times 10^{-5}\left[\frac{\dot{M}}{1M_\odot {\rm y}^{-1}}\right]\left[\frac{M}{M_\odot}\right]^{13/6} \\ &- 8.7\times 10^{-9}\left[\frac{M}{M_\odot}\right] T_9^8\ . \end{aligned}\tag{10}$$

2.4 Angular momentum balance of the star

The angular momentum of the star is changed by two torques induced by accretion and by gravitational radiation if the r-mode oscillation is unstable.

The accretion torque is estimated as $\dot{M}j_a$, where j_a is the specific angular momentum of the accreted matter:

$$j_a = R\sqrt{\frac{2}{7}\frac{GM}{R}}\ . \tag{11}$$

Here the ADAF solution by Narayan & Yi (1994) is employed to evaluate the rotational velocity of the accreted matter at the stellar surface.

As for the gravitational radiation torque, we use the simple formula used in Lindblom et al. (1998) which also incorporates the damping effect by viscosity. This corresponds to the assumption that the r-mode quickly grows to saturate in the non-linear regime of its amplitude and the angular momentum of the mode is a substantial fraction of the stellar angular momentum.

Let us introduce the rate δ of the angular momentum of the mode to the stellar angular momentum. By using the moment of inertia of the star I and the stellar rotational angular frequency Ω, the torque due to gravitational wave emission can be written as ${\tau_{\rm r}}^{-1}I\Omega\delta$. The factor δ is related to the dimensionless amplitude of the oscillation of the r-mode 'κ', which is used in the simplified discussion of the non-linear evolution of the r-mode by Owen et al. (1998), as $\delta \sim \kappa^2$.

The time scale of the r-mode instability $\tau_{\rm r}$ is then defined as,

$${\tau_{\rm r}}^{-1} = {\tau_{\rm gr}}^{-1} + {\tau_{\rm s}}^{-1} + {\tau_{\rm b}}^{-1} \tag{12}$$

where $\tau_{\rm gr}$ is the energy dissipation time scale by the gravitational radiation, $\tau_{\rm s}$ is that of the shear viscosity due to neutron-neutron collision and $\tau_{\rm b}$ is that of the bulk viscosity due to the lag of the $\beta-$reaction and the stellar oscillation. The negative value of $\tau_{\rm r}$ signals the onset of the instability. We set ${\tau_{\rm r}}^{-1}$ to be zero when this value is positive, i.e. for damping oscillations.

The time scale of gravitational wave dissipation is obtained as,

$$\begin{aligned}{\tau_{\rm gr}}^{-1} &= -\frac{32\pi G}{c^{2l+3}}\cdot\frac{(l-1)^{2l}}{[(2l+1)!!]^2}\left(\frac{l+2}{l+1}\right)^{2l+2} \\ &\quad\times\Omega^{2l+2}\rho_c R^{2l+3}\int_0^1 \tilde{\rho}(x)x^{2l+2}dx. \end{aligned}\tag{13}$$

Here $\tilde{\rho}$ is the density distribution normalized by the central density ρ_c. The central density is related to the stellar mass M for $N=1$ polytropes as,

$$\rho_c = 3.29\bar{\rho} = 3.29\times\rho_{\rm nuc}\times 0.7\frac{M}{M_\odot}\ . \tag{14}$$

Thus for $l=m=2$ modes for which the instability is the strongest,

$${\tau_{\rm gr}}^{-1} = -3.1\times 10^{-8}\left[\frac{f}{100{\rm Hz}}\right]^6\left[\frac{M}{M_\odot}\right]. \tag{15}$$

Here f is the rotational frequency of the star ($f=\Omega/2\pi$).

The damping time scales due to shear and bulk viscosities are,

$$\begin{aligned}{\tau_{\rm s}}^{-1} &= (l-1)(2l+1)\int_0^R \eta r^{2l}dr\Big/\int_0^R \rho r^{2l+2}dr \\ &= 5\times 347\rho_c^{9/4}R^5 10^{-18}T_9^{-2}\int_0^1 \bar{\rho}(x)^{9/4}x^4dx \\ &\quad\Big/\left(\rho_c R^7\int_0^1 \bar{\rho}(x)x^6dx\right) \\ &= 2.0\times 10^{-9}\left(\frac{M}{M_\odot}\right)^{5/4} {T_9}^{-2}, \end{aligned}\tag{16}$$

and,

$$\begin{aligned}{\tau_{\rm b}}^{-1} &= (6.99\times 10^8)^{-1}\left(\frac{2\pi f}{\pi G\bar{\rho}}\right)^2 T_9^6 \\ &= 1.4\times 10^{-11}\left[\frac{f}{100{\rm Hz}}\right]^2\left[\frac{M}{M_\odot}\right]^{-1} T_9^6. \end{aligned}\tag{17}$$

The equation for the evolution of the rotational frequency is written as,

$$\begin{aligned}\frac{d}{dt}\left(\left[\frac{M}{M_\odot}\right]\left[\frac{f}{100{\rm Hz}}\right]\right) &= {\tau_r}^{-1}\left[\frac{M}{M_\odot}\right]\left[\frac{f}{100{\rm Hz}}\right]\delta \\ + \ 2.0\times 10^{-6}&\left[\frac{\dot{M}}{1M_\odot{\rm y}^{-1}}\right]\left[\frac{M}{M_\odot}\right]^{1/2}\ . \end{aligned}\tag{18}$$

[¶] Here heating of the star by the r-mode oscillation, which is considered in Levin (1999), is not taken into account. We confirmed, however, that an inclusion of the r-mode heating term is negligible in the situation of our interest.

2.5 Terminal values of the temperature and the rotational frequency of nearly constant mass models

It is important to note that the thermal balance as well as the angular momentum balance determines the approximate terminal values of the temperature and the rotational frequency of the star, if the accretion rate is fixed and the mass is regarded as nearly constant in the evolution.

By setting the right hand side of Eq. (10) to be zero, the temperature can be solved as,

$$T_9 = 3.0 \left[\frac{\dot{M}}{1M_\odot \mathrm{y}^{-1}}\right]^{1/8} \left[\frac{M}{M_\odot}\right]^{7/48} . \tag{19}$$

Thus for the star with mass $M \sim 1.4M_\odot$ and the accretion rate with $\dot{M} \sim 1M_\odot \mathrm{y}^{-1}$, the (uniform) temperature will be about 3.5×10^9K, which is near the most susceptible temperature for the r-mode instability to grow (Lindblom et al. 1998).

The equation of the angular momentum balance gives the terminal value of the rotational frequency of the star by setting the right hand side of Eq.(18) to be zero. If we omit the contributions of viscous damping whose effect is smallest near at the temperature obtained above, we get the expression for the terminal rotational frequency as : ‖

$$\left[\frac{f}{100\mathrm{Hz}}\right] = 1.3 \left[\frac{\dot{M}}{1M_\odot \mathrm{y}^{-1}}\right]^{1/7} \left[\frac{M}{M_\odot}\right]^{-3/14} \delta^{-1/7}. \tag{20}$$

2.6 Evolutionary curves

Given the mass accretion rate and the initial values of the stellar mass, temperature and frequency, the evolution of the corresponding quantities can be followed by integrating Eqs. (1), (10) and (18). We have computed several evolutionary sequences by choosing different sets for the initial values.

We will show how the evolution depends on the initial values. In Fig. 1 the rotational frequency is plotted against the mass of the star which serves as the 'time' variable because the mass accretion rate is assumed to be constant. With the mass accretion rate fixed, solutions with different values of the initial frequency converges to a certain value rather quickly compared with the mass accretion time scale. This behaviour is expected from the rapid temperature saturation seen in Fig. 2 by swift establishment of the thermal balance between heating and cooling. It should be pointed out that the omission of the r-mode instability would lead to very rapid increase of the rotational frequency and that the star would reach its mass-shedding limit before the accretion induced collapse occurs.

The mass accretion rate dependence of the frequency evolution is shown in Fig. 3. As seen in Fig. 2 the temperature of the star quickly saturates at the limiting value which depends on the mass accretion rate. Since the strength of



Figure 1. Evolutionary curves of the rotational frequency $f/100$Hz for different values of the initial rotational frequency, i.e. $f/100\mathrm{Hz} = 0.1, 0.5, 1$ and 3, are plotted against the stellar mass which can be regarded as the 'time' variable. Model parameters are as follows: the initial mass $M = 1.4M_\odot$, the initial temperature $T_9 = 10^{-3}$, the mass accretion rate $\dot{M} = 1M_\odot \mathrm{y}^{-1}$. Numbers attached to the curves are the values of the lapse of time in units of second. Also plotted is the curve of the model in which the r-mode instability is not included (dashed curve).



Figure 2. Evolutionary curves of the temperature $T_9 = T/10^9$(K) for different values of the mass accretion rate, i.e. $\dot{M}/(1M_\odot \mathrm{y}^{-1}) = 1$ (solid curve), 10^{-1} (dotted curve), 10^{-2} (dashed curve), are plotted against the stellar mass. Model parameters are as follows: the initial mass $M = 1.4M_\odot$, the initial temperature $T_9 = 10^{-3}$, and the initial rotational frequency $f/100\mathrm{Hz} = 0.5$.

‖ After the submission, we find a paper on an effect of a magnetic field to the r-mode instability(Rezzolla et al. 2000). Presence of the magnetic field may weaken the instability significantly. If this is the case with the situation of our interest here, the terminal frequency will be much higher.



Figure 3. The rotational frequency is plotted against the stellar mass. Three lines correspond to the three temperature lines in Fig. 2 ($\dot{M}/(1M_{\odot}\mathrm{y}^{-1}) = 1$ (solid curve), 10^{-1} (dotted curve), 10^{-2} (dashed curve)). Numbers attached to these lines are the values of the lapse of time in units of second.

the r-mode instability depends on the temperature as well as the stellar rotational frequency, the rotational frequency also quickly saturates. As the time scale of the angular momentum change is longer than that of the temperature change, the frequency saturation is reached slightly later compared with the temperature saturation.

3 DISCUSSIONS

3.1 Formation of slowly rotating black holes

We have shown that the r-mode instability sets a severe limit on the rotational frequency of the hypercritically accreting neutron stars proposed by Bethe et al. (1999). As the mass accretion rate dependence of the rotational frequency (Eq. (20)) is rather weak, the rotational frequency of the star is well below its mass-shedding limit. Even if the mass accretion rate rises to $\dot{M} = 10^2 M_{\odot}\mathrm{y}^{-1}$, the rotational frequency increases only to 200Hz. Thus the gravitational collapse of neutron stars due to mass accretion to low mass black holes in these systems will proceed almost spherically and black holes formed just after the collapse will rotate so slowly that the rotational parameter is not large either, i.e. $q = cJ/GM^2 \sim 0.01$ for $1.7M_{\odot}$.

3.2 Observational possibility of gravitational waves from the hypercritically accreting neutron stars

The amount of gravitational radiation from the system can be evaluated as follows. The process discussed in this paper is similar to that studied by Wagoner (1984). A neutron star in an X-ray binary system emits gravitational wave so as to balance the accretion torque with the gravitational radiation torque of the f-mode ('Wagoner star'). For our models the f-mode must be replaced by the r-mode and the accretion rate is considerably different, i.e. increased. The standard formula of the luminosity of gravitational radiation can be expressed as (Shapiro & Teukolsky 1983),

$$\frac{c^3}{G}\omega^2 h^2 D^2 \sim 2\pi f \dot{M} j_a \ , \tag{21}$$

where ω, h, D are the eigenfrequency of the mode, the dimensionless strain of the gravitational field and the distance from the source, respectively. Using the terminal rotational frequency f obtained above and the relation of the eigenfrequency and the angular velocity for $l = m = 2$ modes,

$$\omega = \frac{4}{3} \cdot 2\pi f \tag{22}$$

we get

$$h \sim 5.6\times 10^{-22} \left[\frac{\dot{M}}{1M_{\odot}\mathrm{y}^{-1}}\right]^{3/7} \left[\frac{M}{M_{\odot}}\right]^{5/14} \left[\frac{D}{1\mathrm{kpc}}\right]^{-1} \delta^{1/14}. \tag{23}$$

As seen from the result obtained in this paper, the rotational frequency becomes almost constant after a short time of accretion. Thus it is rather easy to detect signals of the gravitational wave by integrating it for an appropriate interval of time.

We here define the Fourier component of the squared dimensionless strain h^2 as $\hat{h}^2$. Then from the Parseval's relation,

$$\hat{h}^2 \cdot \sqrt{\Delta\omega} \sim h^2 \cdot \sqrt{\frac{4\pi}{\omega} N_{\mathrm{cycle}}}, \tag{24}$$

where N_{cycle} is the number of cycles of the oscillation in the interval of observation, Δt_{obs}. Frequency width $\Delta\omega$ and Δt_{obs} are related by the uncertainty principle as,

$$\Delta\omega \cdot \Delta t_{\mathrm{obs}} \geq \frac{1}{2}. \tag{25}$$

As $\Delta t_{\mathrm{obs}} = \frac{4\pi}{\omega} N_{\mathrm{cycle}}$, the r.m.s. of the Fourier component of the strain is,

$$\hat{h} \leq \frac{\sqrt{\Delta t_{\mathrm{obs}}}}{2^{1/4}} h. \tag{26}$$

The event rate $\mathcal{R}$ of the hypercritical accretion is quite uncertain. The maximum value is expected to be $\sim 10^{-4}\mathrm{y}^{-1}\mathrm{galaxy}^{-1}$, which is the birth rate of the close black hole–neutron star binary (Bethe & Brown 1998). In order to detect N_0 events per year, we have to observe the region with the distance as follows:

$$D = (3N_0/4\pi\mathcal{R}n_{\mathrm{gal}})^{1/3} \ , \tag{27}$$

where $n_{\mathrm{gal}} \sim 0.01\mathrm{Mpc}^{-3}$ (Efstathiou et al. 1988). Here n_{gal} is the number density of the massive galaxies ($\sim 10^{11}M_{\odot}$).

Then Fourier component of the strain is,

$$\hat{h}(\mathrm{Hz}^{-1/2}) \lesssim 1 \times 10^{-23} \cdot \delta^{1/14} \left[\frac{\dot{M}}{1M_{\odot}\mathrm{y}^{-1}}\right]^{3/7} \times \left[\frac{M}{M_{\odot}}\right]^{5/14} \left[\frac{\mathcal{R}}{10^{-4}\mathrm{galaxy}^{-1}\mathrm{y}^{-1}}\right]^{1/3} N_0^{-1/3} \left[\frac{t_{\mathrm{obs}}}{1\ \mathrm{month}}\right]^{1/2} \tag{28}$$

Comparing this with the expected sensitivity of the *Laser Interferometer Gravitational-Wave Observatory* (LIGO), the integration of one month is sufficient for the detection by the LIGO II facility (see Shoemaker 2000 for the sensitivity curves of the LIGO). As the dependence of the strain on $\mathcal{R}$ is weak, the LIGO II can see the signal by the integration of one month, even if $\mathcal{R}$ is reduced by two orders of magnitude. Typical frequency of the source may be around $100 \sim 200$Hz.

3.3 Uncertainties in our analysis

In this investigation we have applied the 'one zone' approximation for simplicity. It means that the variables such as the stellar temperature or the rotational frequency are represented by some 'averaged' values throughout the whole star. It premises that the viscosity works very efficiently compared to the time scale of the evolution of the system, typically about one year. The adjustment time scale of shear viscosity to smooth out the differential flow can be estimated for the neutron-neutron scattering dominant matter (Lindblom et al. 1998) as follows:

$$\tau_{\rm adj} \lesssim \frac{R^2\bar{\rho}}{\eta} \sim \frac{R^2 T^2}{347\bar{\rho}^{5/4}}. \qquad (29)$$

Assuming the stellar radius $R = 10$km and the averaged density $\bar{\rho} = 3 \times 10^{14}{\rm g/cm}^3$, this amounts to 0.8 y for $T = 10^8$K. The rotation law of the star can deviate from the uniform rotation if this time scale is comparable or longer than the spin-up time. In that case, behaviour of (classical) r-modes is uncertain and their characteristics may be drastically changed. We here *assume* that they can be approximated by some kind of average of the rotational angular velocity. The validity of this assumption should be confirmed elsewhere.

It has been pointed out that in the hypercritical accretion process, explosive outflows driven by neutrino heating (Fryer et al. 1998) or jet formation (Armitage & Livio 1999) may significantly reduce the mass attachment onto the neutron star. If it is the case, it is more probable that double neutron star binary systems will be formed through this channel. In such accretion inefficient situations, the mass accretion rate $\dot{M}_{\rm acc}$ is lower than the mass inflow rate $\dot{M}_{\rm in}$ from the envelope of the companion star. If the accretion inefficient flows behave similarly as the accretion efficient flows, evolution of the stars may be approximated by our models with lower $\dot{M}$ and with a shorter duration, for the same value of $\dot{M}_{\rm in}$. This results in smaller terminal rotational frequencies of the star. Thus neutron stars which survive after mass accretion in this process would have even lower rotational frequencies.[⋆⋆]

It is a surprise that the possibility of observation of gravitational waves from the systems seems unexpectedly large. This might encourage the ongoing detection experiments of gravitational waves. However we have to be careful in concluding the significance of the hypercritically accreting neutron stars as sources of gravitational waves. Our analysis here might be too simplified and naive to get conclusive answers to the observational possibility. Some simplified assumptions may influence the result significantly. For instance, more realistic behaviour of the accretion flow and the accreted matter may drastically change the efficiency of the accretion torque. Angular momentum re-distribution in the realistic star may modify the r-mode characteristics significantly, which may affect the strength of the instability. Thus the uncertainty included in our assumptions must be investigated more intensively in the future.

ACKNOWLEDGMENT

We would like to express our gratitude to Dr. N. Andersson, the referee, for his suggestion that the detectability of the gravitational wave signals may be rather high.

REFERENCES

Andersson N., 1998, ApJ, 502, 708

Andersson N., Kokkotas K., Stergioulas N., 1999, ApJ, 516, 307

Armitage P. J., Livio M., 1999, ApJ submitted (astro-ph/9906028)

Bethe H. A., Brown G. E., 1998, ApJ, 506, 780

Bethe H. A., Brown G. E., Lee C.-H., 1999, ApJ submitted (astro-ph/9909132)

Brown G. E., 1995, ApJ, 440, 270

Chevalier R. A., 1993, ApJ, 411, L33

Chevalier R. A., 1996, ApJ, 459, 322

Efstathiou G., Ellis R. S., Peterson B., 1988, MNRAS, 232, 431

Friedman J. L., Morsink S. M., 1998, ApJ, 502, 714

Fryer C. L., Benz W., Herant M., 1996, ApJ, 460, 801

Levin Y., 1999, ApJ, 517, 328

Lindblom L., Owen B. J., Morsink S. M., 1998, Phys. Rev. Lett., 80, 4843

Narayan R., Yi I., 1994, ApJ, 428, L13

Owen B. J., Lindblom L., Cutler C., Schutz B. F., Vecchio A., Andersson N., 1998, Phys. Rev. D58, 084020

Rezzolla L., Lamb F. K., Shapiro S. L., 2000, ApJ, 531, L139

Shapiro S. L., Teukolsky S., 1983, *Black Holes, White Dwarfs and Neutron Stars*, Wiley, New York

Shoemaker D., 2000, in *Matters of Gravity, No.15* (http://gravity.phys.psu.edu/mog/mog15/node14.html)

Thorne K. S., 1980, Rev. Mod. Phys, 52, 299

Wagoner R. V., 1984, ApJ, 278, 345

[⋆⋆] It is possible that the characteristics of the accretion flow change drastically and that the accreted matter carries less angular momentum than the case considered here. Of course this will lead to the even more reduction of the terminal rotational frequency.